

## **VORMS DIROFILARIASIS INCREASING EMERGING ZONOSIS IN MONTENEGRO**

*Bogdanka Andric<sup>1</sup>, Snezana Dragas<sup>2</sup> & Bogdan Pajovic<sup>3</sup>*

*<sup>1,2</sup>Professor Clinic for Infectious Disease, Clinic Center of Montenegro*

*Medical School – University of Montenegro, Montenegro*

*<sup>3</sup>Medical faculty University of Montenegro, Podgorica, Krusevac bb, Montenegro*

### **ABSTRACT**

#### **Summary**

*Dirofilariasis belongs to the group of parasitic vector-borne zoonoses. These systemic helminthiases have been widespread distributed in the world, with over 50 million registered cases. Two main filarial species (spp.) have adapted to canine, feline and human hosts: *Dirofilaria repens* and *Dirofilaria immitis*. Human dirofilariases caused inflammatory changes in lymph nodes, lung, irregular and prolonged fever. Different species of infected animals and humans are reservoirs of microfilariae, for blood-sucking arthropod (mosquitoes, flies, ticks). In the vector phase (6 – 10 days), through the process of metamorphosis microfilariae become invasive. In the next phases, feeding arthropods are donors of the microfilariae to sensitive individuals of the humans. The geographical distribution and infectiveness of these vector-borne parasites undergone modifications influenced by global environment and climate changes. Mediterranean area is an endemic region for filarial diseases. Montenegro is a small country in the Balkan region. First three cases of autochthonous dirofilariasis were registered in 2014 / 2015. In Montenegro, there are not data of dirofilariasis in veterinary pathology. In our cases, identification of causes based on surgical extirpation of granulomas and morphologically identification the worms has been performed in three cases.*

**KEYWORDS:** *Dirofilariasis, Diagnosis, Epidemiological, Clinical Aspects*

---

#### **Article History**

**Received: 06 Nov 2018 | Revised: 26 Nov 2018 | Accepted: 01 Dec 2018**

---